

Prothrombin (G724) polyclonal antibody

Catalog: BS2552

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Hemostasis following tissue injury involves the deployment of essential plasma procoagulants (Prothrombin and Factors X, IX, V and VIII), which mediate a blood coagulation cascade that leads to the formation of insoluble Fibrin clots and the promotion of platelet aggregation. Proteolytic cleavage of Prothrombin (Factor II) at residue 44 leads to formation of Thrombin in the first step of the coagulation cascade. Thrombin cleaves bonds between Arg and Gly and activates Factors V, VII, VIII and XIII in complex with thrombomodulin and Protein C. Thrombin maintains vascular integrity during development and postnatal life and coordinates connective tissue proteins by stimulating fibroblast procollagen production.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 70 kDa

Swiss-Prot:

P00734

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

IHC: 1:50~1:200

IF: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Prothrombin (G724) polyclonal antibody detects endogenous levels of Prothrombin protein.

DATA:



Immunohistochemistry (IHC) analyzes of Prothrombin (G724) pAb in paraffin-embedded human breast carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151