

PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



TACC3 (T813) Peptide

Cat No.: BS2580P

Background

TACC1 (transforming acidic coiled coil gene 1) is one of three TACC family members, which are thought to be involved in breast tumorigenesis. TACC1 is located on 8p11 chromosomal region that is amplified in approximately 15% of all breast tumor samples. The short arm of chromosome 8 also contains FGFR1 whose expression is enhanced in most breast cancer tumors. TACC family members, TACC1, TACC2, and TACC3, map very closely to the corresponding FGFR1, FGFR2, FGFR3 genes on chromosomes 4,8, and 10. Subsequently, since they are phylogenetically related, it is proposed that TACC and FGFR have similar roles in cell growth and differentiation. Also, TACC1 contains a conserved C-terminal region as in the Drosophila homolog, D-TACC. It has been shown that D-TACC is necessary for normal spindle function, and the mammalian TACC proteins appears to interact with centrosomes and microtubules in a similar manner.

Swiss-Prot

Q9Y6A5

Applications

Blocking

Specificity

This peptide can be used with studies using BS2580 TACC3 (T813) pAb.

Purification & Purity

Synthetic peptide TACC3 (T813). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

1660 South Highway 100, Suite 500 St. Louis Park, MN

55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co, Ltd.

No 9, weidi road Qixia District Nanjing, 210046,

P, R.China.

Email: info@biogot.com

Tel: +86-025-68037686

Fax: +86-025-68035151