## PRODUCT DATA SHEET



## **Bioworld Technology CO., Ltd.**

# EDG-2 (W45) Peptide

Cat No.: BS2695P

## **Background**

The EDG (endothelial differentiation gene) family of G protein-coupled receptors consists of eight family members that bind lysophospholipid (LPL) mediators, including sphingosine-1-phosphate (SPP) and lysophosphatidic acid (LPA). EDG-1, EDG-3, EDG-5 (also designated H218 and AGR16) and EDG-8 bind SPP with high affinity. EDG-6 is a low affinity receptor for SPP. LPA preferentially binds to EDG-2, EDG-4 and EDG-7. The EDG receptors couple to multiple G proteins to signal through Ras, MAP kinase, Rho, Phospholipase C or other tyrosine kinases, which lead to cell survival, growth, migration and differentiation. EDG-1 signals through Gi proteins to activate Akt and is expressed in glioma cells. EDG-2 is expressed in brain, especially in white matter tract regions, while EDG-3 is expressed in cardiovascular tissue and in cerebellum. EDG-4 is highly expressed on leukocytes and brain, and EDG-5 has wide tissue distribution, including cardiovascular tissue and brain. Expressed in lymphoid and hematopoietic tissues and in lung, EDG-6 signals through Gi/o proteins, which activate growth related pathways.

## **Swiss-Prot**

Q92633

## **Applications**

**Blocking** 

#### **Specificity**

This peptide can be used with studies using BS2695 EDG-2 (W45) pAb.

### **Purification & Purity**

Synthetic peptide EDG-2 (W45). (Note: the amino acid sequence is proprietary). The purity is > 98%.

#### **Product**

1 mg/ml in DI water.

#### **Storage & Stability**

Store at  $4\,\mathrm{C}$  short term. Aliquot and store at  $-20\,\mathrm{C}$  long term. Avoid freeze-thaw cycles.

#### **Research Use**

For research use only, not for use in diagnostic procedure.