

## PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



### EDG-7 (L148) Peptide

Cat No.: BS2698P

#### Background

The EDG (endothelial differentiation gene) family of G protein-coupled receptors consists of eight family members that bind lysophospholipid (LPL) mediators, including sphingosine-1-phosphate (SPP) and lysophosphatidic acid (LPA). EDG-1, EDG-3, EDG-5 (also designated H218 and AGR16) and EDG-8 bind SPP with high affinity. EDG-6 is a low affinity receptor for SPP. LPA preferentially binds to EDG-2, EDG-4 and EDG-7. The EDG receptors couple to multiple G proteins to signal through Ras, MAP kinase, Rho, Phospholipase C or other tyrosine kinases, which lead to cell survival, growth, migration and differentiation. EDG-1 signals through Gi proteins to activate Akt and is expressed in glioma cells. EDG-2 is expressed in brain, especially in white matter tract regions, while EDG-3 is expressed in cardiovascular tissue and in cerebellum. EDG-4 is highly expressed on leukocytes and brain, and EDG-5 has wide tissue distribution, including cardiovascular tissue and brain.

#### Swiss-Prot

Q9UBY5

#### Applications

#### Blocking

#### Specificity

This peptide can be used with studies using BS2698 EDG-7 (L148) pAb.

#### Purification & Purity

Synthetic peptide EDG-7 (L148). (Note: the amino acid sequence is proprietary). The purity is > 98%.

#### Product

1 mg/ml in DI water.

#### Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

#### Research Use

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

1660 South Highway 100, Suite 500 St. Louis Park, MN  
55416, USA.

Email: [info@bioworld.com](mailto:info@bioworld.com)

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co, Ltd.

No 9, weidi road Qixia District Nanjing, 210046,  
P, R.China.

Email: [info@biogot.com](mailto:info@biogot.com)

Tel: +86-025-68037686 Fax: +86-025-68035151