

VPAC1 (V371) polyclonal antibody

Catalog: BS3211

Host: Rabbit

Reactivity: Human, Mouse, Rat

Background:

The vasoactive intestinal peptide (VIP) and the pituitary adenylate cyclase activating polypeptide (PACAP) belong to a superfamily of peptide hormones that include glucagon, secretin and growth hormone releasing hormone. The effects of VIP and PACAP are mediated by three G-protein coupled receptors, VPAC1, VPAC2 and the PACAP receptor (also designated PAC1-R). The VPAC receptors have equal affinities for VIP and PACAP, which stimulate the activation of adenylyl cyclase. Both VPAC1, a 47 kDa protein, and VPAC2, a 65 kDa protein, are abundantly expressed in brain and T cells, where they modulate neuronal differentiation and T cell activation, respectively. The PACAP receptor is a seven transmembrane protein that produces at least eight isoforms by alternative splicing. Each isoform is associated with a specific signaling pathway and a specific expression pattern. The PACAP receptor, which is thought to play an integral role in brain development, preferentially binds PACAP in order to stimulate a cAMP-protein kinase A signaling pathway.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 52 kDa

Swiss-Prot:

P32241

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im-

munogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

IF: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

VPAC1 (V371) polyclonal antibody detects endogenous levels of VPAC1 protein.

DATA:



Immunohistochemistry (IHC) analyzes of VPAC1 (V371) pAb in paraffin-embedded human tonsil carcinoma tissue at 1:50, showing cytoplasmic staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151