

PAR4 (R68) polyclonal antibody

Catalog: BS3238

Host: Rabbit

Reactivity: Human

BackGround:

Thrombin receptor (also designated protease-activated receptor-1 or PAR-1), PAR-2, PAR-3 and PAR-4 compose a distinct class of G protein-coupled receptors activated by proteolysis. Cleavage of these receptors by proteases occurs within the amino-terminal extracellular domain. Thrombin, a serine protease involved in platelet aggregation and blood coagulation, activates the thrombin receptor, resulting in elevated intracellular calcium levels in platelets. Thrombin also cleaves PAR-3 in vitro, suggesting that PAR-3 may be involved in thrombosis or mitogenesis. Thrombin receptor and PAR-4 appear to account for most thrombin signaling in platelets. Activation of PAR-2 in vitro is induced by trypsin, suggesting that PAR-2 is not an alternative thrombin receptor. Cytokines including TNF- α and IL-1 β increase PAR-2 expression, indicating PAR-2 involvement in the acute inflammatory response.

Product:

1 mg/ml in Phosphate buffered saline (PBS) with 15 mM sodium azide, approx. pH 7.2.

Molecular Weight:

~ 41 kDa

Swiss-Prot:

Q96RI0

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IF: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

PAR-4 (R68) polyclonal antibody detects endogenous levels of PAR-4 protein.

DATA:



Western blot (WB) analysis of PAR4 (R68) pAb at 1:500 dilution

Lane1:3T3-L1 whole cell lysate(40ug)

Lane2:PC12 whole cell lysate(40ug)

Lane3:K562 whole cell lysate(40ug)

Lane4:Panc1 whole cell lysate(40ug)

Lane5:H1792 whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151