

IL-8 (V68) polyclonal antibody

Catalog: BS3479

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

IL-8 has been shown to function as a potent neutrophil chemostatic and activating peptide and is an important mediator of inflammatory diseases. Two distinct human IL-8 receptors, designated IL-8RA and IL-8RB, have been characterized. Both are expressed at a high level on neutrophils, and to a lesser extent on monocytes and myeloid cell lines. In addition, the IL-8RA subunit is expressed in T cells such as the Jurkat cell line. Both IL-8Rs are members of the seven-transmembrane domain rhodopsin superfamily of receptors and as such, couple G proteins for signal transduction. The two receptors share 77% amino acid identity.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 11 kDa

Swiss-Prot:

P10145

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


IHC: 1:50~1:200

Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

IL-8 (V68) polyclonal antibody detects endogenous levels of IL-8 protein.

DATA:

Immunohistochemistry (IHC) analyzes of IL-8 (V68) pAb in paraffin-embedded human lung carcinoma tissue at 1:100.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151