

PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



IL-5R α (K68) Peptide

Cat No.: BS3560P

Background

Interleukin 5 (IL-5) is a soluble T cell-derived factor, also known as T cellreplacing factor (TRF), that induces B cell and eosinophil growth and differentiation. IL-5 exerts its biological activity through the IL-5 receptor (IL-5R), which is composed of two chains: an α chain that binds IL-5 with low affinity and a β chain that does not bind IL-5, but together with the IL-5 α chain, constitutes the high affinity IL-5 receptor. The cytoplasmic domain of both the α and β chains is essential for signal transduction. Specifically, the membrane-proximal proline-rich sequence of the cytoplasmic domain of the IL-5R receptor α subunit, IL-5R α , is critical for the IL-5 induced proliferative response, expression of nuclear proto-oncogenes and tyrosine phosphorylation of cellular proteins, such as JAK1 and JAK2.

Swiss-Prot

Q01344

Applications

Blocking

Specificity

This peptide can be used with studies using BS3560 IL-5R α (K68) pAb.

Purification & Purity

Synthetic peptide IL-5R α (K68). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.