PRODUCT DATA SHEET



Bioworld Technology CO., Ltd.

p38 (H174) Peptide

Cat No.: BS3566P

Background

MAP (mitogen-activated protein) kinases play a significant role in many biological processes, including cell adhesion and spreading, cell differentiation and apoptosis. p38 α , p38 β and p38 γ , also known as MAPK14, MAPK11 and MAPK12, respectively, each contain one protein kinase domain and belong to the MAP kinase family. Expressed in different areas throughout the body with common expression patterns in heart, p38 proteins use magnesium as a cofactor to catalyze the ATP-dependent phosphorylation of target proteins. Via their catalytic activity, p38 α , p38 β and p38 γ are involved in a variety of events throughout the cell, including signal transduction pathways, cytokine production and cell proliferation and differentiation.

Swiss-Prot

016539

Applications

Blocking

Specificity

This peptide can be used with studies using BS3566 p38 (H174) pAb.

Purification & Purity

Synthetic peptide p38 (H174). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at $4\,\mathrm{C}$ short term. Aliquot and store at $-20\,\mathrm{C}$ long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.