#### PRODUCT DATA SHEET



## **Bioworld Technology CO., Ltd.**

# GABAA Rδ (D181) Peptide

Cat No.: BS3863P

## **Background**

GAD-65 and GAD-67, glutamate decarboxylases, function to catalyze the production of GABA (gamma-aminobutyric acid). In the central nervous system GABA functions as the main inhibitory transmitter by increasing a Cl- conductance that inhibits neuronal firing. GABA has been shown to activate both ionotropic (GABAA) and metabotropic (GABAB) receptors as well as a third class of receptors called GABAC. Both GABAA and GABAC are ligand-gated ion channels, however, they are structurally and functionally distinct. Members of the GABAA receptor family include GABAA Rα1-6, GABAA R β1-3, GABAA Rγ1-3, GABAA Rδ, GABAA Rε, GABAA Rρ1 and GABAA Rp2. The GABAB family is composed of GABAB R1α and GABAB R1β. GABA transporters have also been identified and include GABA T-1, GABA T-2 and GABA T-3 (also designated GAT-1, -2, and -3). The GABA transporters function to terminate GABA action.

#### **Swiss-Prot**

O14764

## **Applications**

**Blocking** 

#### **Specificity**

This peptide can be used with studies using BS3863 GABAA  $R\delta$  (D181) pAb.

## **Purification & Purity**

Synthetic peptide GABAA R $\delta$  (D181). (Note: the amino acid sequence is proprietary). The purity is > 98%.

#### **Product**

1 mg/ml in DI water.

### **Storage & Stability**

Store at  $4\,\mathrm{C}$  short term. Aliquot and store at -20  $\mathrm{C}$  long term. Avoid freeze-thaw cycles.

#### **Research Use**

For research use only, not for use in diagnostic procedure.