PRODUCT DATA SHEET



Bioworld Technology CO., Ltd.

NMDA_E4 (P706) Peptide

Cat No.: BS3868P

Background

Glutamate receptors mediate most excitatory neurotransmission in the brain and play an important role in neural plasticity, neural development and neurodegeneration. Ionotropic glutamate receptors are categorized into NMDA receptors and kainate/AMPA receptors, both of which contain glutamate-gated, cation-specific ion channels. Kainate/AMPA receptors are co-localized with NMDA receptors in many synapses and consist of seven structurally related subunits designated GluR-1 to -7. The kainate/AMPA receptors are primarily responsible for the fast excitatory neuro-transmission by glutamate, whereas the NMDA receptors exhibit slow kinetsis of Ca2+ ions and a high permeability for Ca2+ ions. The NMDA receptors consist of five subunits: ϵ 1, 2, 3, 4 and one ζ subunit. The ζ subunit is expressed throughout the brainstem whereas the four epsilon subunits display limited distribution.

Swiss-Prot

O15399

Applications

Blocking

Specificity

This peptide can be used with studies using BS3868 NMDAε4 (P706) pAb.

Purification & Purity

Synthetic peptide NMDA ϵ 4 (P706). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at $4\,\mathrm{C}$ short term. Aliquot and store at $-20\,\mathrm{C}$ long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.