

PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



p-ATF2 (T71) Peptide

Cat No.: BS4018P

Background

The transcription factor ATF-2 (also called CRE-BP1) binds to both AP-1 and CRE DNA response elements and is a member of the ATF/CREB family of leucine zipper proteins. ATF-2 interacts with a variety of viral oncoproteins and cellular tumor suppressors and is a target of the SAPK/JNK and p38 MAP kinase signaling pathways. Various forms of cellular stress, including genotoxic agents, inflammatory cytokines and UV irradiation, stimulate the transcriptional activity of ATF-2. Cellular stress activates ATF-2 by phosphorylation of Thr69 and Thr71. Both SAPK and p38 MAPK have been shown to phosphorylate ATF-2 at these sites in vitro and in cells transfected with ATF-2.

Swiss-Prot

P15336

Applications

Blocking

Specificity

This peptide can be used with studies using BS4018 p-ATF2 (T71) pAb.

Purification & Purity

Synthetic peptide p-ATF2 (T71). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.