Bioworld Technology CO., Ltd.



p-Cdk1/Cdc2 (T161) Peptide

Cat No.: BS4033P

Background

Cdc2, an evolutionarily conserved serine/threonine-specific protein kinase, is essential in the cell cycle transition from G2 to M phase. Cdc2 is regulated by association with B-type cyclins and by reversible phosophorylation. Cyclin B binding facilitates the phosphorylation of Cdc2 p34 on three regulatory sites: threonine 14, tyrosine 15, and threonine 161. In higher eukaryotes, Cdc2 is negatively regulated by phosphorylation of two residues located in the ATP-binding site, Thr 14 and Tyr 15. Cdc2 is positively regulated by the cyclin-dependent phosphorylation of Thr 161. Both phosphorylation and de- phosphorylation at Thr 161 are required for progression through the cell cycle.

Swiss-Prot

P06493

Applications

Blocking

Specificity

This peptide can be used with studies using BS4033 p-Cdk1/Cdc2 (T161) pAb.

Purification & Purity

Synthetic peptide p-Cdk1/Cdc2 (T161). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at $4 \,^{\circ}{\rm C}$ short term. Aliquot and store at $-20 \,^{\circ}{\rm C}$ long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.