

## PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



### p-ICAM-1 (Y512) Peptide

Cat No.: BS4096P

#### Background

ICAM1 is a 85-110 kDa single chain type 1 integral membrane glycoprotein with an extracellular domain of five immunoglobulin superfamily repeats, a transmembrane region and a cytoplasmic domain. It shares considerable amino acid sequence homology with ICAM3 and with ICAM2. ICAM1 is expressed by activated endothelial cells. It is detected on cells of many other lineages (e.g. epithelial cells, fibroblasts, chondrocytes, B lymphocytes, T lymphocytes (low), monocytes, macrophages, dendritic cells and neutrophils, with lower levels that increase in inflammation. Also, ICAM1 is detected in some carcinoma and melanoma cells. Soluble ICAM1 is detectable in the plasma and is elevated in patients with various inflammatory syndromes. It is the receptor for rhinoviruses and malaria.

#### Swiss-Prot

P05362

#### Applications

Blocking

#### Specificity

This peptide can be used with studies using BS4096 p-ICAM-1 (Y512) pAb.

#### Purification & Purity

Synthetic peptide p-ICAM-1 (Y512). (Note: the amino acid sequence is proprietary). The purity is > 98%.

#### Product

1 mg/ml in DI water.

#### Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

#### Research Use

For research use only, not for use in diagnostic procedure.