# **Bioworld Technology CO., Ltd.**



# p-PR (S190) Peptide

Cat No.: BS4158P

# Background

The effects of progesterone are mediated by two functionally different isoforms of the progesterone receptor, PR-A and PR-B, which are transcribed from distinct, estrogen inducible promoters within a single copy of the PR gene. The PR-A and PR-B proteins are 90 kDa and 118 kDa respectively; the first 164 amino acids of PR-B are absent in PR-A. Progesterone bound PR-A and PR-B have different transcription activation properties. Specifically, PR-B functions as a transcriptional activator in most cell and promoter contexts, while PR-A is transcriptionally inactive and functions as a strong ligand dependent transdominant repressor of steroid hormone receptor transcriptiona activity. An inhibitory domain (ID), which maps to the amino terminus of the receptor, exists within both PR isoforms.

**Swiss-Prot** 

## P06401

**Applications** 

Blocking

### Specificity

This peptide can be used with studies using BS4158 p-PR (S190) pAb.

#### **Purification & Purity**

Synthetic peptide p-PR (S190). (Note: the amino acid sequence is proprietary). The purity is > 98%.

#### Product

1 mg/ml in DI water.

**Storage & Stability** 

Store at  $4 \,^{\circ}{\rm C}$  short term. Aliquot and store at  $-20 \,^{\circ}{\rm C}$  long term. Avoid freeze-thaw cycles.

#### **Research Use**

For research use only, not for use in diagnostic procedure.