

PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



p-Stat3 (S727) Peptide

Cat No.: BS4180P

Background

Membrane receptor signaling by various ligands, including interferons and growth hormones such as EGF, induces activation of JAK kinases which then leads to tyrosine phosphorylation of the various Stat transcription factors. Stat1 and Stat2 are induced by IFN- α and form a heterodimer which is part of the ISGF3 transcription factor complex. Although early reports indicate Stat3 activation by EGF and IL-6, it has been shown that Stat3 β appears to be activated by both while Stat3 α is activated by EGF, but not by IL-6. Highest expression of Stat4 is seen in testis and myeloid cells. IL-12 has been identified as an activator of Stat4. Stat5 has been shown to be activated by Prolactin and by IL-3. Stat6 is involved in IL-4 activated signaling pathways.

Swiss-Prot

P40763

Applications

Blocking

Specificity

This peptide can be used with studies using BS4180 p-Stat3 (S727) pAb.

Purification & Purity

Synthetic peptide p-Stat3 (S727). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.
1660 South Highway 100, Suite 500 St. Louis Park, MN
55416, USA. Email: info@bioworld.com
Tel: 6123263284 Fax: 6122933841

Bioworld technology, co, Ltd.
No 9, weidi road Qixia District Nanjing, 210046,
P, R.China. Email: info@biogot.com
Tel: +86-025-68037686 Fax: +86-025-68035151