

## PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



### p-PLC $\beta$ 3 (S1105) Peptide

Cat No.: BS4461P

#### Background

A total of eight mammalian PLC isozymes have been described (PLC  $\beta$ 1, PLC  $\beta$ 2, PLC  $\beta$ 3, PLC  $\beta$ 4, PLC  $\gamma$ 1, PLC  $\gamma$ 2, PLC  $\delta$ 1 and PLC  $\delta$ 2) with molecular weights ranging from 85 to 150 kDa. The  $\gamma$ -type enzymes are unique in that they contain SH2 and SH3 domains. Moreover, the two  $\gamma$ -type enzymes, but not the  $\beta$  and  $\delta$  isozymes, are subject to activation by a number of protein tyrosine kinases which associate with their SH2 domains and induce their activation by phosphorylation. In contrast, activation of PLC  $\beta$ 1, PLC  $\beta$ 2 and PLC  $\beta$ 3 is mediated by the  $\alpha$  subunits of the Gq class of heterotrimeric G proteins and by certain  $\beta\gamma$  G protein subunits. The regulatory mechanisms for PLC  $\delta$ 1 and PLC  $\delta$ 2 are as yet not resolved.

#### Swiss-Prot

Q01970

#### Applications

Blocking

#### Specificity

This peptide can be used with studies using BS4461 p-PLC  $\beta$ 3 (S1105) pAb.

#### Purification & Purity

Synthetic peptide p-PLC  $\beta$ 3 (S1105). (Note: the amino acid sequence is proprietary). The purity is > 98%.

#### Product

1 mg/ml in DI water.

#### Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

#### Research Use

For research use only, not for use in diagnostic procedure.