

## CD226 Recombinant Rabbit mAb

BackGround:				1:20		
Involved in intercellular adhesion, lymphocyte signaling,				Storage&Stability:		
cytotoxicity and lymphokine secretion mediated by cyto-				Store at $4{\rm C}$ short term. Aliquot and store at -20 ${\rm C}$ long		
toxic T-lymphocyte (CTL) and NK cell. Cell surface re-				term. Avoid freeze-thaw cycles.		
ceptor for NECTIN2. Upon ligand binding, stimulates				Isotype:		
T-cell proliferation and cytokine production, including				IgG		
that of IL2, IL5, IL10, IL13, and IFNG. Competes with				DATA:		
VRIG for NECT	IN2-binding.					mouse spleen
<b>Product:</b> Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4),			180- 130- 95- 72- 55- ₹ □ CD226 43- 34- 26-			
						0.15M NaCl, 40% Glycerol, 0.01% sodium azide and
0.05% BSA. Stable for 12 months from date of receipt.						
Molecular Weight:						
55-65 kDa						17-
Swiss-Prot:			Western blot detection of CD226 in mouse spleen cell lysates using			
Q8K4F0			CD226 antibody(1:1000 diluted).			
Purification&Purity:			Note:			
Affinity Purification			For research use only, not for use in diagnostic procedure.			
Applications:						
VB: 1:1000 	IHC: 1:200 	ICC/IF: 1	:20 IP:			