

BS46527

Host:

Rabbit

Catalog:

BackGround:

Tissue Factor Pathway Inhibitor Recombinant Rabbit mAb

Reactivity:

Purification&Purity:

Human

This gene encodes a Kunitz-type serine protease inhibitor	Affinity Purification
that regulates the tissue factor (TF)-dependent pathway of	Applications:
blood coagulation. The coagulation process initiates with	WB: 1:1000
the formation of a factor VIIa-TF complex, which prote-	Storage&Stability:
olytically activates additional proteases (factors IX and	Store at 4 ${}^\circ\!\!{\rm C}$ short term. Aliquot and store at -20 ${}^\circ\!\!{\rm C}$ long
X) and ultimately leads to the formation of a fibrin clot.	term. Avoid freeze-thaw cycles.
The product of this gene inhibits the activated factor X	Isotype:
and VIIa-TF proteases in an autoregulatory loop. Inhibi-	IgG
tion of the encoded protein restores hemostasis in animal	DATA:
models of hemophilia. This gene encodes multiple protein	a 12
isoforms that differ in their inhibitory activity, specificity	1800- 130-
and cellular localization. [provided by RefSeq, Jul 2016]	95- 72-
Product:	43- TFPI
Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4),	34-
0.15M NaCl, 40%Glycerol, 0.01% sodium azide and	26-
0.05% BSA. Stable for 12 months from date of receipt.	17-
Molecular Weight:	Western blot detection of TFPI in A549,MCF-7 cell lysates using TFPI
40-55 kDa	antibody(1:1000 diluted).
Swiss-Prot:	Note:
P10646	For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc. Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. Email: info@bioworlde.com Tel: 6123263284 6122933841 Fax:

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: info@biogot.com 0086-025-68037686 Tel: 0086-025-68035151 Fax: