

Nova1 Recombinant Rabbit mAb

BackGround:WB: 1:1000-1:2000 IHC: 1:50-1:200This gene encodes a neuron-specific RNA-binding protein, a member of the Nova family of paraneoplastic disease antigens, that is recognized and inhibited by paraneoplastic antibodies. These antibodies are found in the sera of patients with paraneoplastic opsoclonus-ataxia, breast cancer, and small cell lung cancer. Alternatively spliced transcripts encoding distinct isoforms have been described. [provided by RefSeq, Jul 2008]Store at 20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Jatuate suing Noval in Rat lung cell lysates using Noval antibody.StoresStoresVestern blot detection of Noval in Rat lung cell lysates using Noval antibody.StoresNote:Prification&Purity:For research use only, not for use in diagnostic procedure.Applications:Applications:	Catalog: BS46759 Host: Rabbit	Reactivity: Human, Mouse, Rat
 b. C. C. A. C. A.	BackGround:	WB: 1:1000-1:2000 IHC: 1:50-1:200
ease antigens, that is recognized and inhibited by parane-oplastic antibodies. These antibodies are found in the sera of patients with paraneoplastic opsoclonus-ataxia, breast cancer, and small cell lung cancer. Alternatively spliced transcripts encoding distinct isoforms have been described. [provided by RefSeq, Jul 2008] Product: Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight: 52 kDa Stwiss-Prot: P51513 Purification&Purity: Affinity Purification	This gene encodes a neuron-specific RNA-binding pro-	Storage&Stability:
oplastic antibodies. These antibodies are found in the sera of patients with paraneoplastic opsoclonus-ataxia, breast cancer, and small cell lung cancer. Alternatively spliced transcripts encoding distinct isoforms have been de- scribed. [provided by RefSeq, Jul 2008]Isotype:Product:DATA:Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight:Moter blot detection of Noval in Rat lung cell lysates using Noval atibody.52 kDaWestern blot detection of Noval in Rat lung cell lysates using Noval antibody.51513Note:Purification&Purity: Affinity PurificationFor research use only, not for use in diagnostic procedure.	tein, a member of the Nova family of paraneoplastic dis-	Store at $4\ensuremath{\mathbb{C}}$ short term. Aliquot and store at -20 $\ensuremath{\mathbb{C}}$ long
Of patients with paraneoplastic opsoclonus-ataxia, breast cancer, and small cell lung cancer. Alternatively spliced transcripts encoding distinct isoforms have been de- scribed. [provided by RefSeq, Jul 2008]JataProduct:Datta:Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Mouse brain 1000000000000000000000000000000000000	ease antigens, that is recognized and inhibited by parane-	term. Avoid freeze-thaw cycles.
Def TA:Cancer, and small cell lung cancer. Alternatively spliced transcripts encoding distinct isoforms have been de- scribed. [provided by RefSeq, Jul 2008]Product:Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Molecular Weight:52 kDaSwiss-Prot:P51513Purification&Purity:Arification&Purity:Arification	oplastic antibodies. These antibodies are found in the sera	Isotype:
transcripts encoding distinct isoforms have been described. [provided by RefSeq, Jul 2008] Product: Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight: 52 kDa Swiss-Prot: P51513 Purification&Purity: Affinity Purification	of patients with paraneoplastic opsoclonus-ataxia, breast	IgG
scribed. [provided by RefSeq, Jul 2008] Product: Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight: 52 kDa Swiss-Prot: P51513 Purification&Purity: Affinity Purification	cancer, and small cell lung cancer. Alternatively spliced	DATA:
Product:Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.43- 43- 43- 43- 44- 44- 45- 44- 46- 47- <br< td=""><td></td><td>180- 130-</td></br<>		180- 130-
Store at -20 °C. Supplied in S0mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight: 52 kDa Swiss-Prot: P51513 Purification&Purity: Affinity Purification	Product:	72-
0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight: 52 kDa Swiss-Prot: P51513 Purification&Purity: Affinity Purification	Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4),	43-
Molecular Weight:17.52 kDaWestern blot detection of Noval in Rat lung cell lysates using Noval antibody.Swiss-Prot:Note:P51513Note:Purification&Purity:For research use only, not for use in diagnostic procedure.Affinity PurificationImage: State	0.15M NaCl, 40%Glycerol, 0.01% sodium azide and	
52 kDaWestern blot detection of Noval in Rat lung cell lysates using Noval antibody.Swiss-Prot:Note:P51513Note:Purification&Purity:For research use only, not for use in diagnostic procedure.Affinity PurificationImage: Comparison of the second	0.05% BSA. Stable for 12 months from date of receipt.	
Swiss-Prot:antibody.P51513Note:Purification&Purity:For research use only, not for use in diagnostic procedure.Affinity Purification	Molecular Weight:	17-
Swiss-Trot. Note: P51513 Note: Purification&Purity: For research use only, not for use in diagnostic procedure. Affinity Purification For research use only, not for use in diagnostic procedure.	52 kDa	Western blot detection of Nova1 in Rat lung cell lysates using Nova1
Purification&Purity: For research use only, not for use in diagnostic procedure. Affinity Purification For research use only, not for use in diagnostic procedure.	Swiss-Prot:	antibody.
Affinity Purification	P51513	Note:
-	Purification&Purity:	For research use only, not for use in diagnostic procedure.
Applications:	Affinity Purification	
	Applications:	

Bioworld technology, co. Ltd. Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. Email: info@biogot.com Tel: 0086-025-68037686 Fax: 0086-025-68035151