

## HMGCL Recombinant Rabbit mAb

BackGround:WB: 1:1000 >br/sIHC: 1:100The protein encoded by this gene belongs to the HMG-CoA lyase family. It is a mitochondrial enzyme that catalyzes the final step of leucine degradation and plays a key role in ketone body formation. Mutations in this gene are associated with HMG-CoA lyase deficiency. Alterna- tively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq. Oct 2009]Stora at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.Product:IgG DATA:Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.IgG DATA:Ablecular Weight:Vestern blot detection of HMGCL in Hela cell lysates using HMGCL antibody(1:1000 diluted).Store at -20 °C.Image Stressen	Catalog: BS47199 Host: Rabbit	Reactivity: Human
HMG-CoA lyase family. It is a mitochondrial enzyme that catalyzes the final step of leucine degradation and plays a key role in ketone body formation. Mutations in this gene are associated with HMG-CoA lyase deficiency. Alterna- tively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq. Oct 2009]Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.Product:Isofype:Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Image: Complexite term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.34 kDaWestern blot detection of HMGCL in Hela cell lysates using HMGCL antibody(1:1000 diluted).935914Note:Purification&Purity: Affinity PurificationFor research use only, not for use in diagnostic procedure.	BackGround:	WB: 1:1000 IHC: 1:100
catalyzes the final step of leucine degradation and plays a key role in ketone body formation. Mutations in this gene are associated with HMG-CoA lyase deficiency. Alterna- tively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2009]term. Avoid freeze-thaw cycles.Product:IgGStore at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Image: Comparison of the second seco	The protein encoded by this gene belongs to the	Storage&Stability:
key role in ketone body formation. Mutations in this gene are associated with HMG-CoA lyase deficiency. Alterna- tively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2009]Isotype: IgGProduct:DATA:Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Molecular Weight: a34 kDaWestern blot detection of HMGCL in Hela cell lysates using HMGCL antibody(1:1000 diluted).Profication&Purity: Affinity PurificationNote:Affinity PurificationFor research use only, not for use in diagnostic procedure.	HMG-CoA lyase family. It is a mitochondrial enzyme that	Store at 4 ${\rm C}$ short term. Aliquot and store at -20 ${\rm C}$ long
are associated with HMG-CoA lyase deficiency. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2009]IgGProduct:DATA:Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Image: Comparison of the stable of	catalyzes the final step of leucine degradation and plays a	term. Avoid freeze-thaw cycles.
tively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Oct 2009]DATA:Product:Image: Image: I	key role in ketone body formation. Mutations in this gene	Isotype:
isoforms have been found for this gene. [provided by RefSeq, Oct 2009] Product: Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight: 34 kDa Swiss-Prot: Pa5914 Purification&Purity: Affinity Purification	are associated with HMG-CoA lyase deficiency. Alterna-	IgG
RefSeq, Oct 2009]Product:Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Molecular Weight:34 kDa34 kDaSwiss-Prot:P35914Purification&Purity:Affinity Purification	tively spliced transcript variants encoding different	DATA:
Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Image: Complex of the solution o		180
0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.34Molecular Weight:3534 kDaWestern blot detection of HMGCL in Hela cell lysates using HMGCL antibody(1:1000 diluted).Swiss-Prot:Note:P35914Note:Purification&Purity:For research use only, not for use in diagnostic procedure.Affinity PurificationHela cell lysates using HMGCL	Product:	72- 55-
0.05% BSA. Stable for 12 months from date of receipt.Image: Constant of the constant	Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4),	43-
0.05% BSA. Stable for 12 months from date of receipt.Molecular Weight:34 kDa34 kDaSwiss-Prot:P35914Purification&Purity:Affinity Purification	0.15M NaCl, 40%Glycerol, 0.01% sodium azide and	
34 kDaWestern blot detection of HMGCL in Hela cell lysates using HMGCL antibody(1:1000 diluted).Swiss-Prot:Note:P35914Note:Purification&Purity:For research use only, not for use in diagnostic procedure.Affinity PurificationImage: Comparison of the state of t	0.05% BSA. Stable for 12 months from date of receipt.	
Swiss-Prot:     antibody(1:1000 diluted).       P35914     Note:       Purification&Purity:     For research use only, not for use in diagnostic procedure.       Affinity Purification     For research use only, not for use in diagnostic procedure.	Molecular Weight:	17-
Swiss flot.     Note:       P35914     Note:       Purification&Purity:     For research use only, not for use in diagnostic procedure.       Affinity Purification     For research use only, not for use in diagnostic procedure.	34 kDa	Western blot detection of HMGCL in Hela cell lysates using HMGCL
Purification&Purity:       For research use only, not for use in diagnostic procedure.         Affinity Purification       For research use only, not for use in diagnostic procedure.	Swiss-Prot:	antibody(1:1000 diluted).
Affinity Purification	P35914	Note:
	Purification&Purity:	For research use only, not for use in diagnostic procedure.
Applications:	Affinity Purification	
	Applications:	

Bioworld Technology, Inc.	
Add:	1660 South Highway 100, Suite 500 St. Louis Park,
	MN 55416,USA.
Email:	info@bioworlde.com
Tel:	6123263284
Fax:	6122933841