

PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



p-4E-BP1 (T46) Peptide

Cat No.: BS4747P

Background

The translation of proteins from eukaryotic mRNA is initiated by the multisubunit complex eIF-4F, which associates with the mRNA 5' cap structure. eIF-4E, a component of eIF-4F, is responsible for binding to the 5' cap structure and for the assembly of the eIF-4F complex. The regulatory protein 4E-BP1, also referred to as PHAS-I, inhibits eIF-4E function. Phosphorylation of 4E-BP1 by S6 kinase p70, MAP kinases or PKCs causes the disassociation of 4E-BP1 from eIF-4E, promoting translation. A protein that is functionally related to 4E-BP1, designated 4E-BP2, also associates with eIF-4E.

Swiss-Prot

Q13541

Applications

Blocking

Specificity

This peptide can be used with studies using BS4747 p-4E-BP1 (T46) pAb.

Purification & Purity

Synthetic peptide p-4E-BP1 (T46). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.
1660 South Highway 100, Suite 500 St. Louis Park, MN
55416, USA. Email: info@bioworlde.com
Tel: 6123263284 Fax: 6122933841

Bioworld technology, co, Ltd.
No 9, weidi road Qixia District Nanjing, 210046,
P, R.China. Email: info@biogot.com
Tel: +86-025-68037686 Fax: +86-025-68035151