

# ZAP70 Recombinant Rabbit mAb

Cata	log:	BS47729
------	------	---------

Host:

Rabbit

Reactivity: Human

# **BackGround:**

This gene encodes an enzyme belonging to the protein tyrosine kinase family, and it plays a role in T-cell development and lymphocyte activation. This enzyme, which is phosphorylated on tyrosine residues upon T-cell antigen receptor (TCR) stimulation, functions in the initial step of TCR-mediated signal transduction in combination with the Src family kinases, Lck and Fyn. This enzyme is also essential for thymocyte development. Mutations in this gene cause selective T-cell defect, a severe combined immunodeficiency disease characterized by a selective absence of CD8-positive T-cells. Two transcript variants that encode different isoforms have been found for this gene. [provided by RefSeq, Jul 2008]

### **Product:**

Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.

**Molecular Weight:** 

## 70 kDa

**Swiss-Prot:** 

P43403

Affinity Purification

**Purification&Purity:** 

**Applications:** 

WB: 1:1000<br />ICC/IF: 1:20-1:50<br />FC: 1:20<br />IP: 1:20

### **Storage&Stability:**

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

**Isotype:** 

IgG

**DATA:** 

# Note:

For research use only, not for use in diagnostic procedure.