

HtrA2/Omi Recombinant Rabbit mAb

| Catalog: | BS47909 | Host: | Rabbit | Reactivity: | Human, Mouse, Rat |
|---|--|-------|---------------|---|-------------------|
| | | | | | |
| BackGround: | | | | Swiss-Prot: | |
| This gene encodes a serine protease. The protein has been | | | | O43464 | |
| localized in the endoplasmic reticulum and interacts with | | | | Purification&Purity: | |
| an alternatively spliced form of mitogen-activated protein | | | | Affinity Purification | |
| kinase 14. The protein has also been localized to the mi- | | | | Applications: | |
| tochondria with release to the cytosol following apoptotic | | | | WB: 1:1000-1:2000 IHC: 1:100 IP: 1:20 | |
| stimulus. The protein is thought to induce apoptosis by | | | apoptosis by | Storage&Stability: | |
| binding the apoptosis inhibitory protein baculoviral IAP | | | uloviral IAP | Store at 4 °C short term. Aliquot and store at -20 °C long | |
| repeat-containing 4. Nuclear localization of this protein | | | this protein | term. Avoid freeze-thaw cycles. | |
| has also been observed. Alternate splicing of this gene results in multiple transcript variants encoding different | | | of this gene | Isotype: | |
| | | | ing different | IgG | |
| isoforms. [p | isoforms. [provided by RefSeq, Mar 2016] | | | DATA: | |
| Product: | | | | | |
| | tore at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), | | | Note: | |
| 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and | | | n azide and | For research use only, not for use in diagnostic procedure. | |

Molecular Weight:

0.05% BSA. Stable for 12 months from date of receipt.

36 kDa

Bioworld Technology, Inc.Add:1660 South Highway 100, Suite 500 St. Louis Park,
MN 55416,USA.Email:info@bioworlde.comTel:6123263284Fax:6122933841