

## PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



### p-PI3K p85 $\beta$ (Y464) Peptide

Cat No.: BS4811P

#### Background

Phosphatidylinositol 3-kinase (PI 3-kinase) is composed of p85 and p110 subunits. p85 lacks PI 3-kinase activity and acts as an adapter, coupling p110 to activated protein tyrosine kinase. Two forms of p85 have been described (p85 $\alpha$  and p85 $\beta$ ), each possessing one SH3 and two SH2 domains. Various p110 isoforms have been identified. p110 $\alpha$  and p110 $\beta$  interact with p85 $\alpha$ , and p110 $\alpha$  has also been shown to interact with p85 $\beta$  in vitro. p110 $\delta$  expression is restricted to white blood cells. It has been shown to bind p85 $\alpha$  and p85 $\beta$ , but it apparently does not phosphorylate these subunits. p110 $\delta$  seems to have the capacity to autophosphorylate. p110 $\gamma$  does not interact with the p85 subunits. It has been shown to be activated by  $\alpha$  and  $\beta\gamma$  heterotrimeric G proteins.

#### Swiss-Prot

O00459

#### Applications

Blocking

#### Specificity

This peptide can be used with studies using BS4811 p-PI3K p85 $\beta$  (Y464) pAb.

#### Purification & Purity

Synthetic peptide p-PI3K p85 $\beta$  (Y464). (Note: the amino acid sequence is proprietary). The purity is > 98%.

#### Product

1 mg/ml in DI water.

#### Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

#### Research Use

For research use only, not for use in diagnostic procedure.