

PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



p-PAK2 (S20) Peptide

Cat No.: BS4857P

Background

Three isoforms of serine/threonine kinases, designated α PAK p68, β PAK p65 and γ PAK p62, have been shown to exhibit a high degree of sequence homology with the *S. cerevisiae* kinase Ste 20, involved in pheromone signaling. The α , β and γ PAK isoforms complex specifically with Rac1 and Cdc42 in their active GTP-bound state, inhibiting their intrinsic GTPase activity leading to their autophosphorylation. There are eight sites of autophosphorylation on γ PAK, including Ser 19, Ser 141 and Thr 402, and phosphorylation of Ser 141 and Thr 402 is correlated with γ PAK activation. Once phosphorylated and their affinity for Rac/Cdc42 reduced, the PAK isoforms disassociate from the complex to seek downstream substrates.

Swiss-Prot

Q13177

Applications

Blocking

Specificity

This peptide can be used with studies using BS4857 p-PAK2 (S20) pAb.

Purification & Purity

Synthetic peptide p-PAK2 (S20). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

1660 South Highway 100, Suite 500 St. Louis Park, MN
55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co, Ltd.

No 9, weidi road Qixia District Nanjing, 210046,
P, R.China.

Email: info@biogot.com

Tel: +86-025-68037686 Fax: +86-025-68035151