

# Bioworld Technology CO., Ltd.

# p-PLC v2 (Y1217) Peptide

Cat No.: BS4866P

# **Background**

Phosphoinositide-specific phospholipase C (PLC) plays a crucial role in the initiation of receptor mediated signal transduction through the generation of the two second messengers, inositol 1,4,5-triphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. There are many mammalian PLC isozymes, including PLC β1, PLC β2, PLC β3, PLC β4, PLCγ1, PLC<sub>γ</sub>2, PLC δ1, PLC δ2 and PLCe. PLCδ exists as four different isoforms. PLC δ1, a calcium signal amplifier, is activated by an atypical GTP-binding protein. In addition, PLC δ1 is an effector for GTP-binding protein transglutaminase II-mediated oxytocin receptor and alB-adrenoreceptor signaling. Mouse PLC δ1 is highly expressed in brain, heart, lung and testis. PLC δ is abnormally accumulated in autopsied brains with Alzheimer's disease (AD), suggesting that it may play a role in the pathology of AD. PLC δ2 is markedly expressed in type II intestinal metaplasia and in the adenocarcinoma. When PLC δ2 is expressed in type I intestinal metaplasia, the metaplasia is generally considered benignant, yet evolves toward neoplastic transformation. Thus, PLC δ2 expression may be a possible marker of gastric malignant transformation.

# **Swiss-Prot**

P16885

#### **Applications**

Blocking

#### **Specificity**

This peptide can be used with studies using BS4866 p-PLC γ2 (Y1217) pAb.

#### **Purification & Purity**

Tel: 6123263284

Synthetic peptide p-PLC  $\gamma$ 2 (Y1217). (Note: the amino acid sequence is proprietary). The purity is > 98%.

#### **Product**

1 mg/ml in DI water.

# **Storage & Stability**

Store at  $4 \,\mathrm{C}$  short term. Aliquot and store at  $-20 \,\mathrm{C}$  long term. Avoid freeze-thaw cycles.

# **Research Use**

For research use only, not for use in diagnostic procedure.