

## Prothrombin Recombinant Rabbit mAb

Catalog: BS48760

Host: Rabbit

Reactivity: Human

## **BackGround:**

This gene encodes the prothrombin protein (also known as coagulation factor II). This protein is proteolytically cleaved in multiple steps to form the activated serine protease thrombin. The activated thrombin enzyme plays an important role in thrombosis and hemostasis by converting fibrinogen to fibrin during blood clot formation, by stimulating platelet aggregation, and by activating additional coagulation factors. Thrombin also plays a role in cell proliferation, tissue repair, and angiogenesis as well as maintaining vascular integrity during development and postnatal life. Peptides derived from the C-terminus of this protein have antimicrobial activity against E. coli and P. aeruginosa. Mutations in this gene lead to various forms of thrombosis and dysprothrombinemia. Rapid increases in cytokine levels following coronavirus infections can dysregulate the coagulation cascade and produce thrombosis, compromised blood supply, and organ failure. [provided by RefSeq, May 2020]

## **Product:**

Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and

Molecular Weight:Illy70 kDa70 kDaSwiss-Prot:anP00734Purification&Purity:byAffinity Purificationdi-Applications:inWB: 1:1000<br />ICC/IF: 1:100<br />FC: 1:100Storage&Stability:Store at 4 °C short term. Aliquot and store at -20 °C long<br/>term. Avoid freeze-thaw cycles.Isotype:usIgGDATA:

0.05% BSA. Stable for 12 months from date of receipt.

## Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.	
Add:	1660 South Highway 100, Suite 500 St. Louis Park,
	MN 55416,USA.
Email:	info@bioworlde.com
Tel:	6123263284
Fax:	6122933841