

## **CD3 epsilon Recombinant Rabbit mAb**

| Catalog: BS48816  | Host: Rabbit | Reactivity: Human                          |
|---|--------------|--|
|   |              |  |
| BackGround:   |              | 23 kDa                                     |
| The protein encoded by this gene is the CD3-epsilon   |              | Swiss-Prot:                                |
| polypeptide, which together with CD3-gamma, -delta and<br>-zeta, and the T-cell receptor alpha/beta and gamma/delta<br>heterodimers, forms the T-cell receptor-CD3 complex.<br>This complex plays an important role in coupling antigen<br>recognition to several intracellular signal-transduction<br>pathways. The genes encoding the epsilon, gamma and<br>delta polypeptides are located in the same cluster on<br>chromosome 11. The epsilon polypeptide plays an essen- |              | P07766                                     |
|   |              | Purification&Purity:                       |
|   |              | Affinity Purification                      |
|   |              | Applications:                              |
|   |              | IHC: 1:20-1:100<br>ICC/IF: 1:100<br>       |
|   |              | Storage&Stability:                         |
|   |              | Store at 4 C short term. Aliquot and store |
|   |              | term. Avoid freeze-thaw cycles.            |

tial role in T-cell development. Defects in this gene cause immunodeficiency. This gene has also been linked to a susceptibility to type I diabetes in women. [provided by RefSeq, Jul 2008]

## **Product:**

Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.

**Molecular Weight:** 

/>IP: 1:20

re at -20 ℃ long cies.

## **Isotype:**

IgG

**DATA:** 

## Note:

For research use only, not for use in diagnostic procedure.