

PPIL1 Recombinant Rabbit mAb

BackGround:QSTDC9This gene is a member of the cyclophilin family of petpid dylprolyl isomerases (PPIases). The cyclophilins are a highly conserved, ubiquitous family, members of which play an important role in protein folding, immunosup pression by cyclosporin A, and infection of HIV-1 virions Based on similarity to other PPIases, this protein could accelerate the folding of proteins and might catalyze the isis isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq, Jul 2008]MB: 1:1000 Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.Froduct:Isotype:Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4; 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt:Note:Noteeular Weight:Note:Is kDaNote:Swiss-Prot:Swiss-Prot:	Catalog: BS49058 Host: Rabbit	Reactivity: Human
dylprolyl isomerases (PPIases). The cyclophilins are a highly conserved, ubiquitous family, members of which play an important role in protein folding, immunosup- pression by cyclosporin A, and infection of HIV-1 virions. Based on similarity to other PPIases, this protein could accelerate the folding of proteins and might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq, Jul 2008]Affinity Purification Applications: WB: 1:1000 br />FC: 1:20Product:Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.Isotype:IgGDATA:Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Note: For research use only, not for use in diagnostic procedure.Molecular Weight:Isotype:18 kDaIsotype:	BackGround:	Q5TDC9
Applications:highly conserved, ubiquitous family, members of which play an important role in protein folding, immunosup- pression by cyclosporin A, and infection of HIV-1 virions. Based on similarity to other PPIases, this protein could accelerate the folding of proteins and might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq, Jul 2008]WB: 1:1000 Store at 4°C short term. Aliquot and store at -20°C long term. Avoid freeze-thaw cycles.Product:Store at -20°C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Note: For research use only, not for use in diagnostic procedure.Note:For research use only, not for use in diagnostic procedure.	This gene is a member of the cyclophilin family of pepti-	Purification&Purity:
play an important role in protein folding, immunosuppression by cyclosporin A, and infection of HIV-1 virions.WB: 1:1000 FC: 1:20Based on similarity to other PPIases, this protein could accelerate the folding of proteins and might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq, Jul 2008]Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.Isotype:Isotype:Outct:Isotype:Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Note:Molecular Weight:Note:18 kDaFor research use only, not for use in diagnostic procedure.	dylprolyl isomerases (PPIases). The cyclophilins are a	Affinity Purification
Storage&Stability:pression by cyclosporin A, and infection of HIV-1 virions.Based on similarity to other PPIases, this protein could accelerate the folding of proteins and might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq, Jul 2008]Storage&Stability: Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.Product:Isotype: IgGStore at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.Note: For research use only, not for use in diagnostic procedure.Molecular Weight: 18 kDaIsotype: Image Stability:	highly conserved, ubiquitous family, members of which	Applications:
 Based on similarity to other PPIases, this protein could accelerate the folding of proteins and might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq, Jul 2008] Product: Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight: 18 kDa 	play an important role in protein folding, immunosup-	WB: 1:1000 FC: 1:20
accelerate the folding of proteins and might catalyze the cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq, Jul 2008] Product: Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight: 18 kDa		Storage&Stability:
cis-trans isomerization of proline imidic peptide bonds in oligopeptides. [provided by RefSeq, Jul 2008] Product: Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight: 18 kDa Isotype: IgG DATA: Note: For research use only, not for use in diagnostic procedure.		Store at 4 $^{\circ}\mathrm{C}$ short term. Aliquot and store at -20 $^{\circ}\mathrm{C}$ long
oligopeptides. [provided by RefSeq, Jul 2008] Product: Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight: 18 kDa		term. Avoid freeze-thaw cycles.
Product:DATA:Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40%Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.DATA:Molecular Weight:Note: For research use only, not for use in diagnostic procedure.18 kDa		Isotype:
Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight: 18 kDa		IgG
 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight: 18 kDa Note: For research use only, not for use in diagnostic procedure.		DATA:
0.05% BSA. Stable for 12 months from date of receipt. Molecular Weight: 18 kDa		
Molecular Weight: 18 kDa	-	Note:
18 kDa	•	For research use only, not for use in diagnostic procedure.
	Molecular Weight:	
Swiss-Prot:	18 kDa	
	Swiss-Prot:	

Bioworld Technology, Inc.Add:1660 South Highway 100, Suite 500 St. Louis Park,
MN 55416,USA.Email:info@bioworlde.comTel:6123263284Fax:6122933841