

Ephrin receptor A2 Recombinant Rabbit mAb

Catal	log:	BS49187
-------	------	---------

Host:

Rabbit

Reactivity: Mouse, Rat

BackGround:

Receptor tyrosine kinase which binds promiscuously membrane-bound ephrin-A family ligands residing on adjacent cells, leading to contact-dependent bidirectional signaling into neighboring cells. The signaling pathway downstream of the receptor is referred to as forward signaling while the signaling pathway downstream of the ephrin ligand is referred to as reverse signaling. Activated by the ligand ephrin-A1/EFNA1 regulates migration, integrin-mediated adhesion, proliferation and differentiation of cells. Regulates cell adhesion and differentiation through DSG1/desmoglein-1 and inhibition of the ERK1/ERK2 signaling pathway.

Product:

Store at -20 °C. Supplied in 50mM Tris-Glycine(pH 7.4), 0.15M NaCl, 40% Glycerol, 0.01% sodium azide and 0.05% BSA. Stable for 12 months from date of receipt.

Molecular Weight:

125 kDa

Swiss-Prot:	
Q03145	
Purification&Purity:	
Affinity Purification	
Applications:	
WB: 1:1000 ICC/IF: 1:100 FC: 1:20 IP:	
1:20	
Storage&Stability:	
Store at 4 ${}^\circ\!\!{\rm C}$ short term. Aliquot and store at -20 ${}^\circ\!\!{\rm C}$ long	
term. Avoid freeze-thaw cycles.	
Isotype:	
I C	

IgG

DATA:

Note:

For research use only, not for use in diagnostic procedure.