Bioworld Technology CO., Ltd.



p-PDGFR-β (Y751) Peptide

Cat No.: BS5030P

Background

Platelet-derived growth factor receptors exhibit tyrosine-protein kinase activity and have been implicated in the control of cell proliferation, survival and migration. PDGF receptors, PDGFR-alpha and PDGFR-beta, have 5 extracellular immunoglobulin-like domains and an intracellular tyrosine kinase domain. Upon binding a PDGF, the receptors form homo-and heterodimers. Dimerization of the receptors results in phosphorylain the complex. More than 10 different tion SH2-domain-containing molecules have been shown to bind to different autophosphorylation sites in the PDGF-alpha and beta receptors. PDGF alpha receptors are expressed in oligodendrocyte progenitor cells and PDGF beta receptors are expressed on neurons.

Swiss-Prot

P09619

Applications

Blocking

Specificity

This peptide can be used with studies using BS5030 p-PDGFR- β (Y751) pAb.

Purification & Purity

Synthetic peptide p-PDGFR- β (Y751). (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at $4 \, \mathbb{C}$ short term. Aliquot and store at $-20 \, \mathbb{C}$ long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.