

HSD11B1 polyclonal antibody

Catalog: BS5588

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Glucocorticoid hormone action in target tissues is modulated by 11 β -hydroxysteroid dehydrogenase (11 β -HSD), which catalyzes the interconversion of hormonally active C11-hydroxylated corticosteroids (cortisol, corticosterone) and their inactive C11-keto metabolites (cortisone, 11-dehydrocorticosterone). At least two isoforms of 11 β -HSD exist: a low-affinity NADP-dependent dehydrogenase/oxoreductase (11 β -HSD1) and a high-affinity NAD-dependent dehydrogenase (11 β -HSD2). The glycosylated 11 β -HSD1 protein activates cortisol from cortisone, which is widely expressed in mammals, and is most highly expressed in the liver. 11 β -HSD1 may play a role in glucose homeostasis and pathogenesis of a number of disorders including insulin resistance and obesity.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 32 kDa

Swiss-Prot:

P28845

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

HSD11B1 polyclonal antibody detects endogenous levels of HSD11B1 protein.

DATA:



Western blot (WB) analysis of HSD11B1 polyclonal antibody at 1:500 dilution

Lane1:Hela cell lysate

Lane2:H9C2 cell lysate

Lane3:PC12 cell lysate

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151