

PRODUCT DATA SHEET

Bioworld Technology CO., Ltd.



CD32-C Peptide

Cat No.: BS5656P

Background

CD32 (also designated Fc γ RII) is a low affinity receptor for the Fc fragment of aggregated IgG. CD32 is responsible for the clearance of immunocomplexes by macrophages and also plays an important role in the regulation of antibody production by B cells. IgG can noncooperatively bind either one or two highly glycosylated CD32 molecules, and this binding delivers a negative signal for B cells. CD32 exists as several isoforms that are produced by alternative splicing of three distinct genes, A, B, and C. These isoforms are designated CD32-A, CD32-B1, CD32-B3, and CD32-C. All isoforms are present on monocytes, placental trophoblasts and endothelial cells. In addition, the CD32-B forms are present on B lymphocytes, and the CD32-A and CD32-C forms are found on neutrophils.

Swiss-Prot

P31995

Applications

Blocking

Specificity

This peptide can be used with studies using BS5656 CD32-C pAb.

Purification & Purity

Synthetic peptide CD32-C. (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.