

SMUG1 polyclonal antibody

Catalog: BS5753

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

The base excision repair (BER) pathway removes incorrect bases (uracil) or damaged bases (3-methyladenine) from chromatin. Each BER enzyme system addresses a specific type of base damage. Uracil-DNA glycosylases, UNG2 and SMUG1 (single-strand selective monofunctional uracil DNA glycosylase) remove uracil from both double- and single-stranded DNA in nucleosomes (chromatin core particle). The uracil-excising enzyme family shares structural and functional conservation with minimal sequence conservation. The human SMUG1 gene maps to chromosome 12q13.13.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 34 kDa

Swiss-Prot:

Q53HV7

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200

ICC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

SMUG1 polyclonal antibody detects endogenous levels of SMUG1 protein.

DATA:



Western blot (WB) analysis of SMUG1 polyclonal antibody at 1:500 dilution

Lane1:Hela cell lysate

Lane2:Raw264.7 cell lysate

Lane3:H9C2 cell lysate

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151