

Menin polyclonal antibody

Catalog: BS5795

Host: Rabbit

Reactivity: Human, Mouse, Rat

Background:

Menin (multiple endocrine neoplasia I, MEN1, MEAI, SCG2) is a nuclear tumor suppressor that is mutated in patients with multiple endocrine neoplasia type I (MEN1). Menin can activate the transcription of differentiation-regulating genes by covalent histone modification. In osteoblasts, the interaction of menin and the TGF β /Smad3 pathway negatively regulates BMP2/Smad1/5- and Runx2-dependent transcription activities leading to inhibition of late-stage differentiation. Menin regulates the expression of IGFBP-2 by influencing the IGFBP-2 promoter. Ectopic overexpression of menin via adenoviruses induces apoptosis in murine embryonic fibroblasts in a Bax/Bak-dependent manner. Two mRNA exist and two variants of the shorter mRNA have alternative splicing that changes the CDS. Five variants where alternative splicing takes place in the 5' UTR have been identified.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 68 kDa

Swiss-Prot:

O00255

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000

IHC: 1:50~1:200


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

Menin polyclonal antibody detects endogenous levels of Menin protein.

DATA:




Western blot (WB) analysis of Menin polyclonal antibody at 1:500 dilution

Lane1:HEK293T cell lysate

Lane2:sp2/0 cell lysate

Lane3:H9C2 cell lysate



Immunohistochemistry (IHC) analyzes of Menin pAb in paraffin-embedded human lung carcinoma tissue at 1:50. showing nucleus staining. Negative control (the right) Using PBS instead of primary antibody, secondary antibody is Goat Anti-Rabbit IgG-biotin followed by avidin-peroxidase.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogol.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151