Bioworld Technology CO., Ltd.



Striatin Peptide

Cat No.: BS5949P

Background

Striatin, SG2NA, and zinedin, the three mammalian members of the striatin family, are multimodular, WD-repeat and calmodulin-binding proteins. Zinedin and SG2NA share with striatin identical protein-protein interaction domains and the same overall domain structure. All three proteins are both cytosolic and membrane-bound and bind calmodulin in the presence of calcium. Striatin is a neuronal, intracellular protein strictly expressed in the somato-dendritic compartment, including spines, subsets of neurons, and is considered as a marker of neuronal polarity. Down-regulation of striatin, which is expressed in a few subsets of neurons, impairs the growth of dendrites as well as rat locomotor activity. Zinedin is mainly expressed in the central nervous system, whereas SG2NA is mainly expressed in the brain and muscle.

Swiss-Prot

O43815

Applications

Blocking

Specificity

This peptide can be used with studies using BS5949 Striatin pAb.

Purification & Purity

Synthetic peptide Striatin. (Note: the amino acid sequence is proprietary). The purity is > 98%.

Product

1 mg/ml in DI water.

Storage & Stability

Store at $4 \ \mathbb{C}$ short term. Aliquot and store at $-20 \ \mathbb{C}$ long term. Avoid freeze-thaw cycles.

Research Use

For research use only, not for use in diagnostic procedure.