

PAK6 polyclonal antibody

Catalog: BS60321

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

The p21(CDKN1A)-activated kinases (PAKs) are serine/threonine protein kinases that bind to activated small GTPases, including Cdc42 and Rac, and influence transcription, cell morphology (cytoskeleton rearrangement), motility, and apoptosis. PAK family members contain an amino-terminal Cdc42/Rac interactive binding (CRIB) domain and a carboxyl-terminal kinase domain. PAK6 protein cotranslocates into the nucleus with androgen receptor, which is a steroid hormone-dependent transcription factor that is important for male sexual differentiation and development, in response to androgen. PAK6 transcripts are present at high levels in brain and testis, with lower levels in multiple tissues including prostate and breast.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 75 kDa

Swiss-Prot:

Q9NQ5

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

PAK6 polyclonal antibody detects endogenous levels of PAK6 protein.

DATA:



Western blot (WB) analysis of PAK6 polyclonal antibody at 1:500 dilution

Lane1:Hela cell lysate

Lane2:Mouse brain tissue lysate

Lane3:Rat brain tissue lysate

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151