

TWIST1 polyclonal antibody

Catalog: **BS60412** Host:

Rabbit

Human, Mouse, Rat Reactivity:

BackGround:

Members of the myogenic determination family are basic helix-loop-helix (bHLH) proteins that can be separated into two classes. Class A proteins include the ubiquitously expressed E-box binding factors E12/E47, ITF2 and HEB (BETA1 or HTF4). Class B proteins such as MyoD, myogenin and NeuroD (BETA2) are transiently expressed and exhibit a much more limited tissue distribution. Class A proteins heterodimerize with class B proteins to activate DNA transcription. Working in opposition to these positively acting factors are a specialized group of proteins that function as dominant negative regulators. Muscle tissue is derived from a subset of cells originating from the embryonic mesoderm. The novel basic helix-loop-helix (bHLH) transcription factor twist is a putative regulator of mesodermal differentiation and myogenesis. Twist is expressed throughout the epithelial somite but not in the myotome. Twist requires dimerization with the E proteins and inhibits myogenic regulatory factors. It has been implicated as regulator of the temporal and spatial formation of myotomes.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 22 kDa

Swiss-Prot:

Q15672

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific im-

munogen and the purity is > 95% (by SDS-PAGE).

Applications: WB: 1:500~1:1000


Storage&Stability:

Store at $4 \, \mathbb{C}$ short term. Aliquot and store at $-20 \, \mathbb{C}$ long term. Avoid freeze-thaw cycles.

Specificity:

TWIST1 polyclonal antibody detects endogenous levels of TWIST1 protein.

DATA:



Western blot (WB) analysis of TWIST1 polyclonal antibody at 1:500 dilution

Lane1:C6 whole cell lysate(40ug) Lane2:AML-12 whole cell lysate(40ug) Lane3:SGC7901 whole cell lysate(40ug) Lane4:HepG2 whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. Email: info@bioworlde.com Tel: 6123263284 6122933841 Fax:

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. **Email:** info@biogot.com Tel: 0086-025-68037686 0086-025-68035151 Fax: