

ZNF7 polyclonal antibody

Catalog: BS61378

Host: Rabbit

Reactivity: Human, Mouse, Rat

BackGround:

Zinc finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc finger proteins contain a Kruppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF7 (Zinc finger protein 7), also known as KOX4 or HF.16, is a 686 amino acid zinc-finger protein that belongs to the Kruppel C2H2-type zinc finger family. Localized to the nucleus, ZFP3 contains fifteen C2H2-type zinc fingers and is thought to play a role in transcriptional regulation.

Product:

Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

~ 77 kDa

Swiss-Prot:

P17097

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB: 1:500~1:1000


Storage&Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:

ZNF7 polyclonal antibody detects endogenous levels of ZNF7 protein.

DATA:



Western blot (WB) analysis of ZNF7 polyclonal antibody at 1:500 dilution

Lane1:Hela whole cell lysate(40ug)

Lane2:H9C2 whole cell lysate(40ug)

Lane3:NIH-3T3 whole cell lysate(40ug)

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151