


## **ZNF541** polyclonal antibody

Catalog:	BS61550	Host:	Rabbit	Reactivity:	Human
BackGrou	nd:			Q9H0D2	
Zinc-finger proteins contain DNA-binding domains and				Purification&Purity:	
have a wide variety of functions, most of which encompass some form of tran- scriptional activation				The antibody was affinity-purifi- by affinity-chromatography usi munogen and the purity is $> 95\%$	
or repression. The majority of zinc-finger proteins contain				Applications:	
a Kr üppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF541 (zinc finger protein 541) is a 1,365 amino acid nuclear protein that acts as a com-				WB: 1:500~1:1000	
				Storage&Stability:	
				Store at 4 °C short term. Aliquot term. Avoid freeze-thaw cycles.	
				Specificity:	
remodeling	ponent of a chromatin remodeling multiprotein complex involved in spermato-			ZNF541 polyclonal antibody de of ZNF541 protein.	
genesis. Existing as three alternatively spliced isoforms, ZNF541 interacts				DATA:	
with both H HDAC1, ar main and fiv zinc fingers chromosom	SPA2 and ad contains one SANT we C2H2-type . The gene encoding Z e	SANT domain, a ELM2 do- oding ZNF541 maps to human of consists of over 63 million		180- 130-	2 3
bases, houses approximately				Western blot (WB) analysis of	of ZNF541 n
1,400 genes and is the genetic home for a number of im- munoglobulin				dilution Lane1:L02 whole cell lysate(40ug)	
	(Ig) superfamily members, a number of ICAMs,		CAMs, the	Lane2:A549 whole cell lysat	e(40ug)
	CEACAM and PSG family			Lane3:SK-OVCAR3 whole cell lysate(40	
and Fc receptors (FcRs).				Note:	
Product:				For research use only, not for use	
Rabbit IgG, 1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2					
Molecular Weight:					
~ 160 kDa					
Swiss-Prot	•				

y-purified from rabbit antiserum hy using epitope-specific im->95% (by SDS-PAGE)

Aliquot and store at -20 °C long ycles.

oody detects endogenous levels



2NF541 polyclonal antibody at 1:500

lysate(40ug)

for use in diagnostic procedure.

## Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA. info@bioworlde.com Email: Tel: 6123263284 Fax: 6122933841

## Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China. **Email:** info@biogot.com Tel: 0086-025-68037686 Fax: 0086-025-68035151