

PRODUCT DATA SHEET

Bioworld Technology,Inc.

[KO Validated] RhoGDI polyclonal antibody

Catalog: BS6256 Host: Rabbit Reactivity: Human, Mouse, Rat

BackGround:

This gene encodes a protein that plays a key role in the regulation of signaling through Rho GTPases. The encoded protein inhibits the disassociation of Rho family members from GDP (guanine diphosphate), thereby maintaining these factors in an inactive state. Activity of this protein is important in a variety of cellular processes, and expression of this gene may be altered in tumors. Mutations in this gene have been found in individuals with nephrotic syndrome, type 8. Alternate splicing results in multiple transcript variants.

Product:

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

23kDa

Swiss-Prot:

P52565

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB,1:500 - 1:2000


Storage&Stability:

Store at $4 \, \mathbb{C}$ short term. Aliquot and store at $-20 \, \mathbb{C}$ long term. Avoid freeze-thaw cycles.

Category:

Polyclonal Antibodies

DATA:



Western blot analysis of extracts from normal and RhoGDI knockout 293T cells, using RhoGDI antibody at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.


Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.



Western blot analysis of extracts of various cell lines, using RhoGDI antibody at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.
br/>Lysates/proteins: 25ug per lane.
br/>Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit .
br/>Exposure time: 3s.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: <u>info@bioworlde.com</u>

Tel: 6123263284 Fax: 6122933841 Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: info@biogot.com
Tel: 0086-025-68037686
Fax: 0086-025-68035151