

Retinoic acid induced 17 Polyclonal Antibody

Catalog: BS65987 Host: Rabbit Reactivity: Hu-
man, Mouse, Rat, Chicken, Dog, Pig, Cow, Horse, Sheep,

Background:

Increases ligand-dependent transcriptional activity of AR and promotes AR sumoylation. The stimulation of AR activity is dependent upon sumoylation.

Tissue specificity: Expressed most abundantly in ovary and, at lower levels, in prostate, spleen and testis. Weak expression, if any, in thymus, small intestine, colon and peripheral blood leukocytes.

Product:

0.01M TBS (pH 7.4) with 1% BSA, 0.03% Proclin300 and 50% Glycerol.

Molecular Weight:

115kDa

Swiss-Prot:

Q9ULJ6

Purification & Purity:

affinity purified by Protein A

Applications:

IHC-P=1:100-500 IHC-F=1:100-500 Flow-Cyt=1 μ g/test


Storage & Stability:

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

Specificity:


Retinoic acid induced 17 Polyclonal Antibody detects endogenous levels of Retinoic acid induced 17 protein.

DATA:




Paraformaldehyde-fixed, paraffin embedded (mouse ovary); Antigen

retrieval by boiling in sodium citrate buffer (pH 6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 °C for 30min; Antibody incubation with (Retinoic) Polyclonal Antibody, Unconjugated at 1:200 overnight at 4 °C,



Paraformaldehyde-fixed, paraffin embedded (rat brain); Antigen retrieval by boiling in sodium citrate buffer (pH 6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 °C for 30min; Antibody incubation with (Retinoic) Polyclonal Antibody, Unconjugated at 1:200 overnight at 4 °C



Paraformaldehyde-fixed, paraffin embedded (rat testis); Antigen retrieval by boiling in sodium citrate buffer (pH 6.0) for 15min; Block endogenous peroxidase by 3% hydrogen peroxide for 20 minutes; Blocking buffer (normal goat serum) at 37 °C for 30min; Antibody incubation with (Retinoic) Polyclonal Antibody, Unconjugated at 1:200 overnight at 4 °C,

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151



PRODUCT DATA SHEET

Bioworld Technology, Inc.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,
MN 55416, USA.

Email: info@bioworld.com

Tel: 6123263284

Fax: 6122933841

Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,
P. R. China.

Email: info@biogot.com

Tel: 0086-025-68037686

Fax: 0086-025-68035151