

# NMDAR2A polyclonal antibody

Catalog: BS70491

Host: Rabbit

Reactivity: Mouse, Rat

## **BackGround:**

This gene encodes a member of the glutamate-gated ion channel protein family. The encoded protein is an N-methyl-D-aspartate (NMDA) receptor subunit. NMDA receptors are both ligand-gated and voltage-dependent, and are involved in long-term potentiation, an activity-dependent increase in the efficiency of synaptic transmission thought to underlie certain kinds of memory and learning. These receptors are permeable to calcium ions, and activation results in a calcium influx into post-synaptic cells, which results in the activation of several signaling cascades. Disruption of this gene is associated with focal epilepsy and speech disorder with or without mental retardation. Alternative splicing results in multiple transcript variants.

**Product:** 

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

**Molecular Weight:** 

## 180KDa

**Swiss-Prot:** 

Q12879

**Purification&Purity:** 

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

#### **Applications:**

WB,1:500 - 1:1000|IF/ICC,1:50 - 1:200


**Storage&Stability:** 

Store at  $4 \,^{\circ}{\rm C}$  short term. Aliquot and store at  $-20 \,^{\circ}{\rm C}$  long term. Avoid freeze-thaw cycles.


#### Category:

**Polyclonal Antibodies** 


### **DATA:**



Western blot analysis of extracts of various cell lines, using NMDAR2A antibody at 1:1000 dilution.<br/>Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>br/>Lysates/proteins: 25ug per lane.<br/>br/>Blocking buffer: 3% nonfat dry milk in TBST.<br/>br/>Detection: ECL Basic Kit .<br/>br/>Exposure time: 180s.



Immunofluorescence analysis of H9C2 cells using NMDAR2A Rabbit pAb at dilution of 1:100. Blue: DAPI for nuclear staining.



Immunofluorescence analysis of L929 cells using NMDAR2A Rabbit pAb at dilution of 1:100. Blue: DAPI for nuclear staining.

#### Note:

For research use only, not for use in diagnostic procedure.

## Bioworld Technology, Inc.

 
 Add:
 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.

 Email:
 info@bioworlde.com

 Tel:
 6123263284

 Fax:
 6122933841

## Bioworld technology, co. Ltd.

 
 Add:
 No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

 Email:
 info@biogot.com

 Tel:
 0086-025-68037686

 Fax:
 0086-025-68035151