

**ALKBH7 polyclonal antibody**

Catalog: BS70810

Host: Rabbit

Reactivity: Human, Mouse, Rat

**BackGround:**

May function as protein hydroxylase; can catalyze auto-hydroxylation at Leu-110 (in vitro, but this activity may be due to the absence of the true substrate. Required to induce programmed necrosis in response to DNA damage caused by cytotoxic alkylating agents. Acts by triggering the collapse of mitochondrial membrane potential and loss of mitochondrial function that leads to energy depletion and cell death. ALKBH7-mediated necrosis is probably required to prevent the accumulation of cells with DNA damage. Does not display DNA demethylase activity. Involved in fatty acid metabolism (By similarity.

**Product:**

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

**Molecular Weight:**

26-30kDa

**Swiss-Prot:**

Q9BT30

**Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

**Applications:**


WB,1:200 - 1:2000

**Storage&Stability:**

Store at 4 °C short term. Aliquot and store at -20 °C long term. Avoid freeze-thaw cycles.

**Category:**

Polyclonal Antibodies

**DATA:**

Western blot analysis of extracts of various cell lines, using ALKBH7 antibody at 1:1000 dilution. Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution. Lysates/proteins: 25ug per lane. Blocking buffer: 3% nonfat dry milk in TBST. Detection: ECL Basic Kit. Exposure time: 90s.

**Note:**

For research use only, not for use in diagnostic procedure.

**Bioworld Technology, Inc.**

Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416, USA.

Email: [info@bioworld.com](mailto:info@bioworld.com)

Tel: 6123263284

Fax: 6122933841

**Bioworld technology, co. Ltd.**

Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.

Email: [info@biogot.com](mailto:info@biogot.com)

Tel: 0086-025-68037686

Fax: 0086-025-68035151