

GABA A Receptor beta 2 (GABRB2) polyclonal antibody

Catalog:	BS7120	Host:	Rabbit	Reactivity: Human, Mouse
BackGround:				munogen and the purity is $> 95\%$ (by SDS-PAGE).
The gamma-aminobutyric acid (GABA) A receptor is a				Applications:
multisubunit chloride channel that mediates the fastest			s the fastest	WB,1:500 - 1:2000
inhibitory synaptic transmission in the central nervous			tral nervous	Storage&Stability:
system. This gene encodes GABA A receptor, beta 2				Store at 4 ${}^\circ\!\!{\rm C}$ short term. Aliquot and store at -20 ${}^\circ\!\!{\rm C}$ long
subunit. It is mapped to chromosome 5q34 in a cluster				term. Avoid freeze-thaw cycles.
comprised of genes encoding alpha 1 and gamma 2 subu-			nma 2 subu-	Category:
nits of the GABA A receptor. Alternative splicing of this			-	Polyclonal Antibodies
gene generates 2 transcript variants, differing by a 114 bp			by a 114 bp	DATA:
insertion.				
Product:				Western blot analysis of extracts of various cell lines, using GABAA
1mg/ml in PBS with 0.02% sodium azide, 50% glycerol,)% glycerol,	Receptor beta 2) Rabbit pAb at 1:1000 dilution. Secondary anti-
pH7.2				body: HRP Goat Anti-Rabbit IgG at 1:10000 dilu-
Molecular Weight: 55KDa Swiss-Prot:				tion. Lysates/proteins: 25ug per lane. Blocking buffer: 3%
				nonfat dry milk in TBST. br/>Detection: ECL Basic Kit . br/>Exposure
				time: 180s.
P47870				Note:
Purification&Purity:				For research use only, not for use in diagnostic procedure.
The antibody was affinity-purified from rabbit antiserum				

Bioworld Technology, Inc.Add:1660 South Highway 100, Suite 500 St. Louis Park,
MN 55416,USA.Email:info@bioworlde.comTel:6123263284Fax:6122933841

by affinity-chromatography using epitope-specific im-