

# PRODUCT DATA SHEET

Bioworld Technology,Inc.

# DMT1/SLC11A2 polyclonal antibody

Catalog: BS71487 Host: Rabbit Reactivity: Human, Mouse, Rat

### **BackGround:**

This gene encodes a member of the solute carrier family 11 protein family. The product of this gene transports divalent metals and is involved in iron absorption. Mutations in this gene are associated with hypochromic microcytic anemia with iron overload. A related solute carrier family 11 protein gene is located on chromosome 2. Multiple transcript variants encoding different isoforms have been found for this gene.

#### **Product:**

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

# **Molecular Weight:**

72KDa

## **Swiss-Prot:**

P49281

## **Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

## **Applications:**

WB,1:1000 - 1:2000|IHC,1:50 - 1:200


## **Storage&Stability:**

Store at  $4\,\mathrm{C}$  short term. Aliquot and store at  $-20\,\mathrm{C}$  long term. Avoid freeze-thaw cycles.


## **Category:**

Polyclonal Antibodies


#### **DATA:**



Western blot analysis of extracts of various cell lines, using DMT1/SLC11A2 Rabbit pAb at 1:1000 dilution.<br/>
secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>
br/>Lysates/proteins: 25ug per lane.<br/>
br/>Blocking buffer: 3% nonfat dry milk in TBST.<br/>
br/>Detection: ECL Basic Kit .<br/>
secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>
secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>
secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>
secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>
secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>
secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>
secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>
secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>
secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>
secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>
secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>
secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>
secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.



Immunohistochemistry of paraffin-embedded rat stomach using DMT1/SLC11A2 Rabbit pAb at dilution of 1:100 .Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.



Immunohistochemistry of paraffin-embedded mouse stomach using DMT1/SLC11A2 Rabbit pAb at dilution of 1:100 .Perform microwave antigen retrieval with 10 mM Tris/EDTA buffer pH 9.0 before commencing with IHC staining protocol.

### Note:

For research use only, not for use in diagnostic procedure.

# Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: <u>info@bioworlde.com</u>

Tel: 6123263284 Fax: 6122933841 Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: <u>info@biogot.com</u>
Tel: 0086-025-68037686
Fax: 0086-025-68035151