

## PRODUCT DATA SHEET

Bioworld Technology,Inc.

# Granulin polyclonal antibody

Catalog: BS7852 Host: Rabbit Reactivity: Human, Mouse

#### **BackGround:**

Granulins are a family of secreted, glycosylated peptides that are cleaved from a single precursor protein with 7.5 repeats of a highly conserved 12-cysteine granulin/epithelin motif. The 88 kDa precursor protein, progranulin, is also called proepithelin and PC cell-derived growth factor. Cleavage of the signal peptide produces mature granulin which can be further cleaved into a variety of active, 6 kDa peptides. These smaller cleavage products are named granulin A, granulin B, granulin C, etc. Epithelins 1 and 2 are synonymous with granulins A and B, respectively. Both the peptides and intact granulin protein regulate cell growth. However, different members of the granulin protein family may act as inhibitors, stimulators, or have dual actions on cell growth. Granulin family members are important in normal development, wound healing, and tumorigenesis.

# **Product:**

1 mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

#### **Molecular Weight:**

88KDa

### **Swiss-Prot:**

P28799

## **Purification&Purity:**

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

# **Applications:**

WB,1:500 - 1:2000|IF/ICC,1:50 - 1:200

## Storage&Stability:


Store at 4 ℃ short term. Aliquot and store at -20 ℃ long

term. Avoid freeze-thaw cycles.


#### **Category:**

Polyclonal Antibodies

#### **DATA:**



Western blot analysis of extracts of A-431 cells, using Granulin antibody at 1:1000 dilution.<br/>Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.<br/>dr/>Lysates/proteins: 25ug per lane.<br/>Blocking buffer: 3% nonfat dry milk in TBST.<br/>Detection: ECL Basic Kit.<br/>Exposure time: 180s.



Immunofluorescence analysis of NIH/3T3 cells using Granulin Rabbit pAb at dilution of 1:200 . Blue: DAPI for nuclear staining.

## Note:

For research use only, not for use in diagnostic procedure.

## Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: <u>info@bioworlde.com</u>

Tel: 6123263284 Fax: 6122933841 Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: <u>info@biogot.com</u>
Tel: 0086-025-68037686
Fax: 0086-025-68035151