

PRODUCT DATA SHEET

Bioworld Technology,Inc.

[KO Validated] RB polyclonal antibody

Catalog: BS79514 Host: Rabbit Reactivity: Human

BackGround:

The protein encoded by this gene is a negative regulator of the cell cycle and was the first tumor suppressor gene found. The encoded protein also stabilizes constitutive heterochromatin to maintain the overall chromatin structure. The active, hypophosphorylated form of the protein binds transcription factor E2F1. Defects in this gene are a cause of childhood cancer retinoblastoma (RB), bladder cancer, and osteogenic sarcoma.

Product:

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

110kDa

Swiss-Prot:

P06400

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB,1:500 - 1:2000|IF/ICC,1:50 - 1:200


Storage&Stability:

Store at $4\,\mathrm{C}$ short term. Aliquot and store at $-20\,\mathrm{C}$ long term. Avoid freeze-thaw cycles.

Modification:

Unmodification

DATA:



Western blot analysis of extracts of various cell lines, using RB antibody at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

br/>Lysates/proteins: 25ug per lane.

Blocking buffer: 3% nonfat dry milk in TBST.


br/>Detection: ECL Basic Kit.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

br/>Blocking buffer: 3% nonfat dry milk in TBST.

Blocking buffer: 3% nonfat dry milk in TBST.

Blocking buffer: 3% nonfat dry milk in TBST.



Western blot analysis of extracts from normal and RB Rabbit pAb knockout HeLa cells, using RB Rabbit pAb antibody at 1:1000 dilution.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

br/>Lysates/proteins: 25ug per lane.

Blocking buffer: 3% nonfat dry milk in TBST.


Detection: ECL Basic Kit.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.

Blocking buffer: 3% nonfat dry milk in TBST.

Detection: ECL Basic Kit.

Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.



Immunofluorescence analysis of Y79 cells using Rb Polyclonal Antibody at dilution of 1:100 . Blue: DAPI for nuclear staining.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.

Add: 1660 South Highway 100, Suite 500 St. Louis Park,

MN 55416,USA.

Email: info@bioworlde.com

Tel: 6123263284 Fax: 6122933841 Bioworld technology, co. Ltd.

Add: No 9, weidi road Qixia District Nanjing, 210046,

P. R. China.

Email: <u>info@biogot.com</u> Tel: 0086-025-68037686 Fax: 0086-025-68035151