ACSS2 polyclonal antibody

Catalog: BS8244 Host: Rabbit Reactivity: Human, Mouse, Rat

BackGround:

This gene encodes a cytosolic enzyme that catalyzes the activation of acetate for use in lipid synthesis and energy generation. The protein acts as a monomer and produces acetyl-CoA from acetate in a reaction that requires ATP. Expression of this gene is regulated by sterol regulatory element-binding proteins, transcription factors that activate genes required for the synthesis of cholesterol and unsaturated fatty acids. Alternative splicing results in multiple transcript variants.

Product:

1mg/ml in PBS with 0.02% sodium azide, 50% glycerol, pH7.2

Molecular Weight:

78kDa

Swiss-Prot:

Q9NR19

Purification&Purity:

The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen and the purity is > 95% (by SDS-PAGE).

Applications:

WB,1:500 - 1:2000|IF/ICC,1:50 - 1:200

Storage&Stability:

Store at 4℃ short term. Aliquot and store at -20℃ long term. Avoid freeze-thaw cycles.

Category:

Polyclonal Antibodies

DATA:

Western blot analysis of extracts of various cell lines, using ACSS2 antibody at 1:1000 dilution.
Secondary antibody: HRP Goat Anti-Rabbit IgG at 1:10000 dilution.
Lysates/proteins: 25ug per lane.
Blocking buffer: 3% nonfat dry milk in TBST.
Detection: ECL Basic Kit .
Exposure time: 5s.

Confocal immunofluorescence analysis of NIH-3T3 cells using ACSS2 Polyclonal Antibody at dilution of 1:200. Blue: DAPI for nuclear staining.

Immunofluorescence analysis of NIH-3T3 cells using ACSS2 Rabbit pAb at dilution of 100 . Blue: DAPI for nuclear staining.

Note:

For research use only, not for use in diagnostic procedure.

Bioworld Technology, Inc.
Add: 1660 South Highway 100, Suite 500 St. Louis Park, MN 55416,USA.
Email: info@bioworlde.com
Tel: 6123263284
Fax: 6122933841

Bioworld technology, co. Ltd.
Add: No 9, weidi road Qixia District Nanjing, 210046, P. R. China.
Email: info@biogot.com
Tel: 0086-025-68037686
Fax: 0086-025-68035151